Brief ASAM Screening Tool

Client Name:	Date:					
City/geographic region: _		_	Phone:			
Were responses provided	by a third party? Yes	□ No	Okay to	o leave V/	′M □ Yes	□ No
Referral source:	□ Accoss/Crisis	Line (ACL) □ Dro	ug Donon	dancy Car	urt	
□ Sell	□ Access/Crisis		ug Depen	deficy co	uit	
☐ CalWorks Case Manage	ement Other:	□ De	cline to s	tate		
Gender Identity: ☐ Male	☐ Female	☐ Transgender (M to	F)	☐ Trans	sgender (F to	M)
☐ Questioning/Unsure	☐ Other	☐ Decline to state		☐ Unkn	own	
If female, are you current	ly pregnant? ☐ Yes ☐ No	0				
Sexual Orientation:						
☐ Heterosexual/Straight ☐ Lesbian		\square Gay	☐ Bisexual			
☐ Questioning/Unsure	☐ Other:	Decline to s	□ Unknown			
Dimension 1: Withdrawa	I/Detox Potential					
	cing any current severe with miting, excessive sweating,					
If YES to 1, make	immediate referral for medi	cal evaluation of need for	acute, ir	patient c	are. Stop Sc	reen.
2. Are you under the	e influence of any substance	s right now?	☐ Yes	□ No	☐ Unknow	n □ N/A
3. If NO, have you us	sed any substances in the la	st 1-3 days?	☐ Yes	□ No	☐ Unknow	n □ N/A
If YES to 2, consid	er Withdrawal Managemen	t. Continue screening.				
4. How does drinkin	g alcohol/using drugs impac	t your daily life or function	oning? Ple	ase descr	ribe:	

Comments:					
S	everity		(Substance Use, Acute In k one of the following lev		otential)
☐ 0: None		☐ 1: Mild	☐ 2: Moderate	☐ 3: Significant	☐ 4: Severe
Fully functioning, n signs of intoxication W/D present.	n or	Mild to moderate intoxication interferes with daily functioning, but does not pose a danger to self/others. Minimal risk of severe W/D.	Intoxication may be severe, but responds to support; not posing a danger to self or others. Moderate risk of severe W/D.	Severe signs/symptoms of intoxication indicate an imminent danger to self/others. Risk of severe but manageable W/D; or W/D is worsening.	Incapacitated, with severe signs/symptoms. Severe W/D presents danger, such as seizures. Continued use poses an imminent threat to life (e.g., liver failure, GI bleeding, or fetal death).
Dimension 2: Bio	omedica	al Conditions and Com	plications (not related to	withdrawal)	
1. Are you	having a	medical emergency?		☐ Yes ☐ No	☐ Unknown ☐ N/A
If YES to	1, make	immediate referral fo	r medical evaluation of n	eed for acute, inpatient c	are. Stop Screen.
2 Doyou h	2240 225	, physical boalth condi	tions or disabilities? If you	dossribo bolow	
•		Unknown \square N/A	tions or disabilities? If yes	, describe below.	
If yes to	#2, do a	ny of these health con	ditions have an impact or	n your daily life or functio	ning?
				☐ Yes ☐ No	☐ Unknown ☐ N/A
If yes, de	escribe: _.				

3.		e any special accommodat 		her?) □ Yes □ No 	☐ Unknown ☐ N/A
Comme	ents:				
		Severity Rating – Dimens			
□ 0: No	one	Please Check ☐ 1: Mild	one of the following level \square 2: Moderate	<i>ls of severity</i> ☐ 3: Significant	☐ 4: Severe
Fully fun	nctioning and cope with any discomfort or	Adequate ability to cope with physical discomfort. Mild to moderate symptoms (such as mild to moderate pain) interfere with daily functioning.	Some difficulty tolerating physical problems. Acute, non-life threatening medical symptoms (such as acute episodes of chronic, distracting pain, or signs of malnutrition or electrolyte imbalance) are present. Serious biomedical problems are neglected.	Poor ability to tolerate and cope with physical problems, and/or general health condition is poor. Serious medical problems neglected during outpatient or IOT services. Severe medical problems (such as severe pain requiring medication, or hard to control Type 1 Diabetes) are present but stable.	The person is incapacitated, with severe medical problem (such as extreme pain, uncontrolled diabetes, GI bleeding, or infection requiring IV antibiotics).
	Are you currer	al/Behavioral/Cognitive (and the control of substitution of substitution) and explain in the control of substitution of substi	cide or hurting yourself?	☐ Yes ☐ No ☐] Unknown □ N/A de or hurt yourself:
2.	-	ntly having thoughts of cau			
	If YES to 2, ask	additional questions to fu	orther assess for Tarasoff.	If found to be a Taraso	ff incident, follow
3.	-	ntly experiencing a behavio	oral health crisis, such as s	severe mental or emotio	nal issues?

ii yes, doc	ument additional information	ı in detaii:		
	r indicated due to client's ans I respond as directed by agen	•		emergency facility,
•	ve a mental health diagnosis? cify:			□ Unknown □ N/A
☐ Yes ☐	4, does your mental health ha No □ Unknown □ N/A cribe:			
Comments:				
Severity	Rating – Dimension 3 (Emotic	onal Behavioral or Cogni	tive (FBC) Conditions or (Complications)
	Please Check	one of the following leve	els of severity	
☐ 0: None	1: Mild	☐ 2: Moderate	☐ 3: Significant	4: Severe
Good impulse control coping skills and subdomains (dangerousness/lethinterference with recovery efforts, soci functioning, self-care ability, course of illness	diagnosed EBC condition that requires intervention, but does not significantly interfere with treatment. Relationships are being	Persistent EBC condition, with symptoms that distract from recovery efforts, but are not an immediate threat to safety and do not prevent independent functioning.	Severe EBC symptomatology, but sufficient control that does not require involuntary confinement. Impulses to harm self/others, but not dangerous in a 24-hr. setting	Severe EBC symptomatology; requires involuntary confinement. Exhibits severe and acute lifethreatening symptoms (e.g., dangerous or impulsive behavior or cognitive functioning) posing imminent danger
				to self/others.

Dimension 4: Readiness to Change 1. Have you been mandated or directed to receive SUD (substance use disorder) treatment? ☐ Yes ☐ No ☐ Unknown ☐ N/A If yes, describe mandate/direction: 2. How ready are you to change your alcohol or drug use now? \Box Unknown \Box N/A ☐ Not ready ☐ Getting Ready ☐ Ready \square In process of ☐ Sustained change making changes made (Maintenance) Comments: Severity Rating – Dimension 4 (Readiness to Change) Please check one of the following levels of severity ☐ 3: Significant □ 0: None ☐ 1: Mild ☐ 2: Moderate ☐ 4: Severe Unable to follow through, Engaged in treatment as Ambivalent of the need Reluctant to agree to Minimal awareness of a proactive, responsible to change. Willing to treatment. Able to need to change. Only little or no awareness of explore need for participant. Committed articulate negative partially able to follow problems, knows very through with treatment to change. treatment and strategies consequences (of little about addiction, to reduce or stop substance use and/or recommendations. sees no connection substance use. May mental health problems) between substance believe it will not be but has low commitment use/consequences. Not difficult to change, or to change. Passively willing to explore change. does not accept a full involved in treatment Unwilling/unable to recovery treatment plan. (variable follow through, follow through with variable attendance) treatment recommendations.

Dimension 5: Relapse/Continued Use Potential

		0	1	2	3	4	5	6	7	8	9	10	likely
Nor	ne						5						Very
	desire/urge to use substances? $\ \square$ Unknown $\ \square$ N/A												
3.	3. On a scale from 0 to 10, with 0 being "none" and 10 being "very likely", how would you describe your												
2.	2. Are you likely to continue to drink or use without treatment? \Box Yes \Box No \Box Unknown \Box N							□ N/A					
1.	Have you drank or used on most days (15 or more) in the last 30 days? \Box Yes \Box No \Box Unknown \Box N/ A									□ N/A			

Comments:									
Severity Rating – Dimension 5 (Relapse, Continued Use, or Continued Problem Potential) Please check one of the following levels of severity									
☐ 0: None	Please ☐ 1: Mild	check one of the following the collection ☐ 2: Moderate	_	Significant ☐ 3: Significant		☐ 4: Severe			
Low or no potential for further substance use problems or has low relapse potential. Goo coping skills in place.	potential. Some risk fair coping and relap		apse elf-	Little recognition and understanding of relissues, poor skills to with relapse.	and Repeated treatment relapse episodes have had little				
Dimension 6: Recovery Environment 1. Is your current living situation unsafe or harmful to your recovery?									
Comments:									
		ng – Dimension 6 (Red check one of the follo							
☐ 0: None	☐ 1: Mild	☐ 2: Moderate		Significant	□ 4	: Severe			
Supportive environment and/or able to cope in environment.	Passive/disinterested social support, but not too distracted by this situation and still able to cope.	Unsupportive environment, but able to cope with clinical structure most of the time.	Unsupple and the	portive environment e client has difficulty , even with clinical	recover friend availated environ Unabone negation environ envi	onment toxic/hostile to very (i.e. many drug-using ds, or drugs are readily able in the home onment, or there are nic lifestyle problems). It is to cope with the tive effects of this onment on recovery (i.e. onment may pose a threat covery).			

Level of Care Inquiry: Do you have an idea about the type of treatment you are interested in? ☐ Yes ☐ No ☐ Unknown ☐ N/A ☐ OTP/MAT ☐ Outpatient ☐ Intensive Outpatient ☐ Residential Other: ☐ Withdrawal Management **Level of Care Disposition:** Recommended Level of Care: ☐ Outpatient ☐ Intensive Outpatient ☐ Residential ☐ OTP/MAT ☐ Withdrawal Management ☐ Urgent/Crisis Actual Level of Care Offered: ☐ Outpatient ☐ Intensive Outpatient ☐ Residential ☐ OTP/MAT ☐ Withdrawal Management ☐ Urgent/Crisis Reason for Discrepancy (if any): ☐ Service not available ☐ Provider judgment ☐ Not Applicable ☐ Client preference ☐ Transportation ☐ Accessibility ☐ Financial ☐ Preferred to wait ☐ Language/Cultural Factors ☐ Environment ☐ Mental Health ☐ Physical Health ☐ Court/Probation ordered ☐ Other _____ ☐ Client on waiting list for indicated level Program referral(s): Printed Name: Signature:

Date: